

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING


Kendon

All Seasons White Oil Insecticide

ACTIVE CONSTITUENT: 825g/L PETROLEUM OIL

FOR THE CONTROL OF SCALE INSECTS ON CITRUS AND GRAPES,
MITES ON POME AND STONE FRUIT TREES, AND CERCOSPORA,
CORDANA, LEAF SPOTS, LEAF SPECKLE ON BANANAS, INSECTS
ON ORNAMENTALS AND FOR USE AS A SPREADER AS PER
THE **DIRECTIONS FOR USE** IN THE
ATTACHED **LABEL LEAFLET**

**IMPORTANT: READ THE ATTACHED
LABEL LEAFLET BEFORE USING
THIS PRODUCT**



5L
10L
20L

AN AUSTRALIAN OWNED COMPANY

Kendon Chemical & MNFG. CO. Pty Ltd. 10 Abbott St, Fairfield, Victoria 3078 Ph 03 9497 2822



**PROTECTION OF WILDLIFE, FISH,
CRUSTACEA AND ENVIRONMENT:**

Do not contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL:

Keep out of reach of children.

Store in the closed, original container in a well ventilated area, as cool as possible.

Do not store for prolonged periods in direct sunlight. Triple or preferably, pressure rinse empty containers before disposal. Add rinsings to spray tank.

Do not dispose of undiluted chemicals on-site. Break, crush and or puncture and bury empty containers in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should be burnt. In the event of spillage, absorb onto sand or chemical absorbent. Remove residue with

detergent and hot water. Dispose of by normal waste disposal methods conforming to local authority codes.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre on 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET:

Additional information is listed in the Material Safety Data Sheet which can be supplied from supplier.

CONDITIONS OF SALE:

This product as supplied is of high grade and believed to be suitable for any purpose for which it is expressly supplied and must be used in accordance with the directions for use given on this label. No responsibility is accepted in respect of this product, save those non-excludable conditions implied by the Trade Practices Act or any applicable State Legislation.

**IN A TRANSPORT
EMERGENCY DIAL 000
POLICE OR FIRE BRIGADE**

**SPECIALIST ADVICE IN
EMERGENCY ONLY
(03) 9497 2822**

APVMA No. 49333/5-20L/0507

BN

DOM

5L



10L



20L



CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENDON ALL SEASONS WHITE OIL INSECTICIDE

ACTIVE CONSTITUENT:
825g/L PETROLEUM OIL

For the control of Scale Insects on Citrus and Grapes,
Mites on Pome and Stone Fruit trees and Cercospora,
Cordana, Leaf Spots, Leaf Speckle on Bananas,
Insects on Ornamentals and for use as a Spreader,
as per the DIRECTIONS FOR USE table.

**IMPORTANT: READ THIS LEAFLET
BEFORE USING THIS PRODUCT**

Kendon Chemical & Mnfg Co Pty Ltd
10 Abbott Street, Fairfield VIC 3078
Telephone: 03 9497 2822

AN AUSTRALIAN OWNED COMPANY, EST. 1950

DIRECTIONS FOR USE
RESTRAINTS: DO NOT spray when shade temperatures exceed 30°C.
DO NOT spray when soil is dry or trees are suffering from moisture stress.

| CROP | PEST | STATE | RATE | CRITICAL COMMENTS |
|---|---------------------------------|----------------------------------|---|--|
| Citrus, except 'Glen Retreat' variety Mandarins | Citrus Leafminer | QLD, NSW, ACT, VIC, TAS, SA & WA | For control of Citrus Leafminer use 500ml / 100L. | During each flush cycle spray every 5 to 14 days. In summer and early autumn, spraying will be needed more frequently than in mid to late autumn. Sprays must be applied thoroughly, ensuring that susceptible leaves are covered on both sides. PRECAUTIONS: if petroleum spray oil is applied more than once from spring to autumn, either alone or in combination with other materials, the total strength of oil used in commercial orchards should not exceed 2L in Queensland and South Australia and 2.5L in NSW and Victoria. DO NOT spray citrus with sprays containing 1L to 1.2L product / 100L water before October or after mid March. DO NOT spray when the soil is dry: trees must not be suffering from moisture stress when sprayed. Ensure that oil-water mixtures held in spray tanks or containers during application are continually agitated or stirred to prevent the oil separating from the water. Petroleum oil sprays greater than 250ml product / 100L water should not be applied for at least one month after spraying with sulphur or lime sulphur, as severe injury may occur, especially if the oil is applied in hot weather. |
| | | | If there is also a need to control other pests eg. Red Scales, then use 500ml - 1L / 100L | |
| | | | | |
| | | | | |
| | Red Scale | VIC, SA & WA | 1.2L / 100L | Apply between mid December and mid March. |
| | | NSW & ACT | 1.3L - 2L / 100L | |
| | | QLD | 1L - 1.7L / 100L | |
| | White Wax Scale, Pink Wax Scale | VIC, WA | 1.2L / 100L | Apply between mid December and mid March. |
| | | NSW & ACT | 1.3L - 2L / 100L | |
| | | QLD | 1L / 100L | Application late November to early December. Another application late January to early February if required. Tank mix with methidathion 50g / 100L. |
| | | | | Application late November to early December. Another application late January to early February, if required. Tank mix with methidathion 50g/100L, carbaryl 70g/100L, promecarb 50g/100L, soda ash 500g-750g/100L or washing soda 500g-750g/100L. |

| CROP | PEST | STATE | RATE | CRITICAL COMMENTS |
|---|---|-----------------------------|--------------------|--|
| Citrus, except 'Glen Retreat' variety Mandarins cont... | Black Scale | TAS & WA | 1.8L - 2.5L / 100L | Application mid December to early January. Apply again in February. |
| | Soft Brown Scale | TAS & WA | 1.8L - 2.5L / 100L | Application January / April. |
| Pome Fruit, Stone Fruit | San Jose Scale, Oyster Shell Scale, Bryobia Mite Eggs, European Red Mite Eggs | NSW, ACT, VIC, TAS, SA & WA | 2L / 100L | Application during dormant period up to green tip (Pome Fruit), or Bud Swell (Stone Fruit). For WA, apply in combination with a recommended chemical treatment for San Jose Scale. |
| | | QLD | 3L / 100L | |
| | Two Spotted Mite | QLD, NSW & ACT | 3L / 100L | Application during dormant period up to green tip (Pome Fruit) or Bud Swell (Stone Fruit). |
| Bananas | Cercospora Leaf Spot | NSW | 4L / Ha | Apply in combination with mancozeb at 2.4kg - 4.5kg / Ha in 200L - 500L of water. Apply at 3-4 week intervals from December until late April. In Nth QLD the intervals may be reduced to 10-14 days in the wet season. |
| | | QLD | 3.5L / Ha | |
| | Cercospora Leaf Speckle, Leaf Spot | NSW & Sth QLD | 12L / Ha | As above in a minimum of 12L of water. Starting in November, early December, apply as a mist at intervals of not more than 3 weeks until April. Spray both sides of leaves. |
| | | NSW & Sth QLD | 5L / Ha | Apply with Propicol 250EC at 200ml - 400ml / Ha. Begin spraying in December and apply 4-5 times at 21 day intervals. If required, use alternative fungicides after this. Spray both sides of leaves. |
| | Cercospora Leaf Spot, Leaf Speckle, Cordana Leaf Spot | Nth QLD | 5L / Ha | Apply with Propicol 250EC at 200ml - 400ml / Ha. Spray from the start of the wet season in December every two weeks for a maximum of 6 applications. If required, use alternative fungicides after this. Spray both sides of leaves. |
| | Scale Insects | QLD, NSW & WA | 2L / 100L | Use when heavy scale populations are present on foliage, fruit or stems. If trees need watering, do not apply. Efficacy is best against young crawler stages. |

| CROP | PEST | STATE | RATE | CRITICAL COMMENTS |
|--|--|-----------------------------|----------------------------------|---|
| Avocado | Scale Insects | QLD, | 1.2L OR 1.6L / 100L | Use when heavy scale populations are present on foliage, fruit or stems. If trees need watering, do not apply. Efficacy is best against young crawler stages. |
| | | NSW & WA | 2L / 100L | |
| Kiwi Fruit | Scale Insects | QLD | 1L - 1.6L / 100L | |
| | | NSW & WA | 2L / 100L | |
| Custard Apple | Scale Insects | QLD | 1L - 1.7L / 100L | |
| | | NSW & WA | 2L / 100L | |
| Mango | Scale Insects | QLD | 1.25L / 100L | |
| | | NSW & WA | 2L / 100L | |
| Grapes | Grapevine Scale | QLD, NSW, ACT, VIC, SA & WA | 3L / 100L | Control is possible if the dormant vines are sprayed in winter, preferably after pruning. |
| Roses | Scale, Aphids, Mites & Mealy Bugs | All | 2L / 100L | Apply in winter after pruning and before bud burst or at other times when pests appear. |
| Shade Trees* and Shrubs, including Conifers, Deciduous, Broadleaf Evergreens and Woody Ornamentals, Ornamental Trees*. Shrubs along city streets, other rights of way, including Deciduous and Broadleaf Evergreens | Aphids, Scales (immature), Spider Mites, Whitefly (immature) | NSW, VIC, SA, WA & TAS | 2L - 4L (Winter) | Winter dormant period as needed for plants that become dormant in winter. |
| | | | 1L - 3L (Summer) | Summer (foliar or cover) as needed. |
| Plants & Ornamental Shrubs | Scale Insects, Mealy Bugs & Spider Mites | | | Spray when pests appear. Repeat after 4 weeks. |
| Daphne | | | 1L / 100L | |

| CROP | PEST | STATE | RATE | CRITICAL COMMENTS |
|---------------|-----------------|-----------------------------|--------------|--|
| As a Spreader | All pests above | QLD, NSW, VIC, TAS, SA & WA | 500ml / 100L | Use in conjunction with parathion, malathion, thiram and ziram, if used at low oil strengths. Spray coverage is improved with the inclusion of Kendon All Seasons White Oil. |

*** Oil removes the glaucous (blue) from some evergreens.**
Always use lower dosage or test spray oil sensitive plants first.
Do not spray Walnut foliage.

| PEST | SPRAYER | VOLUME OF SPRAY (L / Ha) | | |
|--------------------|---------------------------------|--------------------------|-------------------|-------------|
| | | 500ml** | 1L** | 2L** |
| Red Scale | Oscillating Boom with Outrigger | | 8000-12000 | |
| | Airblast with Tower | | 9000-14000 | |
| | Rotary Atomiser | | 8000-12000 | |
| Young Soft Scales | Oscillating Boom with Outrigger | | 6000-10000 | 6000-10000 |
| | Airblast with Tower | | 6000-10000 | 6000-10000 |
| | Rotary Atomiser | | 6000-8000 | 6000-8000 |
| Citrus Red Mite | Oscillating Boom with Outrigger | 7000-10000 | 5000-8000 | |
| | Airblast with Tower | 7000-10000 | 5000-8000 | |
| | Rotary Atomiser | 7000-8000 | 4500-6500 | |
| Rust Mites | Oscillating Boom with Outrigger | 7000-10000 | 5000-8000 | |
| | Airblast with Tower | 7000-10000 | 5000-8000 | |
| | Rotary Atomiser | 7000-8000 | 4500-8000 | |
| Citrus Leaf Miner | Oscillating Boom with Outrigger | 3000-4000 | | |
| | Airblast with Tower | 3000-4000 | | |
| | Rotary Atomiser | 3000-4000 | | |
| Young Mealy Bugs* | Oscillating Boom with Outrigger | | 10000 | 10000-12000 |
| | Airblast with Tower | | 10000 | 10000-14000 |
| | Rotary Atomiser | | 8000 | 8000-10000 |
| Greenhouse Thrips* | Oscillating Boom with Outrigger | | 10000-12000 | |
| | Airblast with Tower | | 10000-12000 | |
| | Rotary Atomiser | | 8000-12000 | |

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION
™Registered Trade Mark of Hoechst Schering Agrevo GmbH, Federal Republic of Germany

WITHHOLDING PERIOD: DO NOT apply later than 1 day before harvest.

GENERAL INSTRUCTIONS

Apply when trees are in good condition 7-10 days after rain or irrigation.

MIXING INSTRUCTIONS

Pour the required amount of water into the mixing tank, then add the Kendon All Seasons White Oil and stir well. When using in combination with other pesticides, mix the pesticide first according to the manufacturers instructions, then add Kendon All Seasons White Oil last. Maintain contnuuous agitation while spraying.

SPRAYING INSTRUCTIONS

Refer to the table for recommended spray volumes. These are based on mature, four metre high orange trees (250 single planted or 500 double planted trees per hectare). Volumes in **bold** typeface mean that the same spray may need to be applied twice in one season. Volumes in normal typeface indicate that one spray per season is generally sufficient. Where a range of values is given, the lower value is suitable for low to medium infestations in the presence of good natural enemy activity. The higher value is for heavy infestations, and may also be suitable for the production of fresh fruit. Application for the control of soft scales, wax scales and mealy bugs must be applied when the youngest stages of the pests predominate.

*Single season applications, or the first of two sprays, for Greenhouse Thrips and Mealy bugs should be made post harvest, initially in late spring / early summer, before new season fruits start to touch.

**Volume of Kendon All Seasons White Oil per 100L of water. Runoff should not be considered as wastage as it is generally better to apply higher volumes with less oil than lower volumes with more oil. Thorough coverage is very important and must include the tops of trees. Maintain continuous agitation while spraying.

PROTECTION OF CROPS, NATIVE AND OTHER NON TARGET PLANTS

DO NOT apply Sertin™ 186EC and Kendon All Seasons White Oil in tank mixes of methazole or ioxynil as crop damage may occur.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Keep out of reach of children.

Store in the closed, original container in a well ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Triple or preferably, pressure rinse empty containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Break, crush, and or puncture and bury empty containers in a local authority landfill. If not available, bury the containers below 500mm, in a disposal pit specifically marked and set up for this purpose, clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

In the event of spillage, absorb onto sand or chemical absorbant. Remove residue with detergent and hot water. Dispose of by normal waste disposal methods, conforming to local authority codes.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 131 126). If swallowed, **DO NOT** induce vomitting. Give a glass of water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS), which can be obtained from the supplier.

CONDITIONS OF SALE

This product as supplied is of high grade and believed to be suitable for any purpose for which it is expressly supplied and must be used in accordance with the directions for use given on this label. No responsibility is accepted in respect of this product, save those non excludable conditions implied by the Trade Practices Act or any applicable State Legislation.

| |
|---|
| GHS Statements Causes serious eye irritation. May be fatal if swallowed and enters airways. |
|---|

APVMA No. 49333/0507